

認知症予防研究成果

Original articles

1. Kitamura K, Nakamura K, Ueno K, Nishiwaki T. Cognitive function is maintained in noninstitutionalized elderly Japanese requiring care with high levels of health-related quality of life. *Environ Health Prev Med.* in press.
2. Nakamura K, Kitamura K, Watanabe Y, Shinoda H, Sato H, Someya T. Rural-urban differences in the prevalence of cognitive impairment in independent community-dwelling elderly residents of Ojiya city, Niigata Prefecture, Japan. *Environ Health Prev Med.* in press.
3. Kitamura K, Watanabe Y, Nakamura K, Sanpei K, Wakasugi M, Yokoseki A, Onodera O, Ikeuchi T, Kuwano R, Momotsu T, Narita I, Endo N. Modifiable factors associated with cognitive impairment in 1143 Japanese outpatients: The Project in Sado for Total Health (PROST). *Dement Geriatr Cogn Dis Extra.* 2016;6:341-9.
4. Watanabe Y, Kitamura K, Nakamura K, Sanpei K, Wakasugi M, Yokoseki A, Onodera O, Ikeuchi T, Kuwano R, Momotsu T, Narita I, Endo N. Elevated C-reactive protein is associated with cognitive decline in outpatients of a general hospital: The Project in Sado for Total Health (PROST). *Dement Geriatr Cogn Dis Extra.* 2016;6:10-9.